

12

EUROPEAN PATENT APPLICATION

21 Application number: **89810155.5**

61 Int. Cl.4: **A 61 K 37/02**

22 Date of filing: **28.02.89**

30 Priority: **04.03.88 GB 8805231**
09.05.88 GB 8810899

43 Date of publication of application:
06.09.89 Bulletin 89/36

64 Designated Contracting States: **ES GR LU**

71 Applicant: **SANDOZ AG**
Lichtstrasse 35
CH-4002 Basel (CH)

72 Inventor: **Feyen, Jean Honoré M.**
Mühlegasse 2
CH-4104 Oberwil (CH)

Trechsel, Ulrich
Bollwerkstrasse 60
CH-4102 Binningen (CH)

The title of the invention has been amended (Guidelines for Examination in the EPO, A-III, 7.3).

54 **Use of an Interleukin-6 molecule in inhibiting bone resorption.**

57 Interleukin-6 molecules are useful in inhibiting bone resorption and increasing bone formation.

EXPRESS MAIL NO. EV545395094US
DOCKET NO. PC32145A

EP 0 331 640 A2

Description

BONE RESORPTION

This invention relates to bone resorption.

Interleukin 6 is a cytokine which has been shown to be produced by fibroblasts and many other cell types. Production of interleukin 6 and its activity on, e.g. bone marrow is described in PCT application PCT/US 87/01611 filed off US Appln No. 06/885905 filed July 15, 1986 (WO 88/00206). Interleukin 6 appears to have multiple target cells. Interleukin 6 turned out to be identical to interferon beta-2, B-cell stimulatory factor 2 (BSF-2), interleukin HP-1 and 26K factor. It is now doubtful whether interleukin-6 can be regarded as an interferon at all in view of its low anti-viral activity. It has been reported that the production of interleukin 6 can be stimulated by interleukin 1 (IL-1) and Tumour Necrosis Factor (TNF α) in certain cells. It is also known that prostaglandin E₂ production, a marker of bone resorption, is stimulated by IL-1 and TNF α .

Despite intensive research with interleukin-6 in particular immunological and hematological areas, there has been no report of effects of interleukin 6 on bone resorption or formation.

We have now found that IL-6 molecules (as defined hereinafter) are useful in inhibiting bone resorption and increasing bone formation. They are thus useful in treating osteoporosis, e.g. acute and/or chronic stages of postmenopausal osteoporosis, including primary and/or secondary osteoporosis and localized osteoporosis, post traumatic osteoporosis, Paget's disease of bone (osteitis deformans), osteolysis due to bone malignancies, e.g. osteogenic sarcoma, osteodystrophy, e.g. renal osteodystrophy, osteomalacia, multiple myeloma of the bone, osteogenesis imperfecta, hypercalcemia, osteomyelitis, e.g. in conjunction with antibiotics and ectopic calcification.

The results below show that IL-6 molecules (as defined hereinafter):

- i) inhibit PGE₂ secretion from bone cell cultures (Test 1)
- ii) inhibit PGE₂ secretion stimulation induced by e.g. two lymphokines, interleukin-1 and tumour necrosis factor, in bone cultures (Test 2),
- iii) increase bone formation (Test 3), and
- iv) decrease bone resorption (Test 4 and Test 5).

The increase in bone formation is of particular interest.

Additionally it is shown that osteoblasts, which also mediate osteoclasts, produce interleukin-6 activity (Tests 6 and 7).

As used hereinafter 1 unit (U) of interleukin-6 activity refers to the amount of IL-6 molecule which causes half-maximal stimulation in the B-cells proliferation assay e.g. using clone B13.29 (which is freely available from the Central Laboratory Netherlands Red Cross, Blood Transfusion Service, P.O.Box 9406, 1006 AK Amsterdam) as described in the specific assay of Landsdorp et al. Curr.Top. Microbiol. Immunol. (1986) 132, 105. and Aarden et al. Eur. J. Immunol. (1987) 17, 1411. 1 x 10⁸ units is about 1 mg of polypeptide.

In brief, cells are suspended in culture medium (Iscove's modified Dulbecco's medium supplemented with 5% (v/v) fetal calf serum, penicillin and streptomycin) and plated (5.10³ cells /200 μ L medium) in microtiter plates. Cells are cultured for 48 hours, in a humidified atmosphere of 5% CO₂ in air at 37°C, the last 6 hours in the presence of 1 microCi of [³H] thymidine and the radioactivity incorporated in the nuclei analysed.

The conditioned medium to be tested is serially diluted in B13.29 cell medium.

Interleukin-6 for which test data is given refers to glycosylated IL-6 produced by CHO cells, if desired purification to remove LPS (lipopolysaccharide).

As used hereinafter Interleukin-1 α = IL-1.

Test 1

Basal production of prostaglandin E₂ in the rat osteosarcoma cell line ROS 17/2.8 is inhibited at a concentration of from about 0.01 to about 100 U/ml IL-6 molecule. In this test cells of the rat osteosarcoma cell line ROS 17/2.8 are cultured in Dulbecco's minimal essential medium and medium F12 (1:1) containing 10% fetal calf serum and 2 mM L-glutamine in a humidified atmosphere of 5% CO₂ in air at 37°C. At confluency, cells are treated with the above concentration of IL-6 molecule. After 24 hours of culture an aliquot of the medium is assayed for prostaglandin E₂ using a standard radioimmunoassay.

Results obtained with interleukin-6 are as follows:-

Concentration of interleukin-6 in U/ml	PGE ₂ Pg (picograms)/ml
0 (Control)	184
0.1	151
1.0	128
10	118

Test 2

Production of prostaglandin E₂ produced by stimulation of mouse calvarial cells by interleukin-1 and tumor necrosis factor is reduced at a concentration of from about 10 to about 100 Units/ml of IL-6 molecule.

In this test organ cultures of mouse calvariae are first prepared:-

Frontal and parietal bones are dissected from neonatal (5 to 6-day-old) mice (strain CD-1), split along the sagittal suture, and cultured in 2 ml of BGJ- medium containing 1 mg/ml bovine serum albumin, penicillin and streptomycin in 35 mm plastic tissue culture wells.

After the preculture period (24 hours), the medium is replaced with BGJ-culture medium supplemented with interleukin-1 at a concentration of 10 units or TNFalpha at a concentration of 10 nanograms/ml and the IL-6 molecule at the above concentration and cultured for another 48 hours. During the entire culture period, the multi-well dishes are shaken on a rocker platform (15 oscillations/min) in a humidified atmosphere of 5% CO₂ in air at 37°C. The calvaria-conditioned medium is harvested and placed at 4°C.

An aliquot of the medium is then assayed for prostaglandin E₂ using a radioimmunoassay.

Results obtained with interleukin-6 are as follows:-

Stimulation by 10 U/ml interleukin-1

Concentration of Interleukin-6 (Units/ml)	Prostaglandin E ₂ pg/ml
0	4800
10	4400
100	500*

(* equivalent to control value without stimulation by interleukin-6).

Stimulation by 10 ng/ml TNFa

Concentration of Interleukin-6 (Units/ml)	Prostaglandin E ₂ pg/ml
0	3300
10	1200
100	800*

(* Control value without stimulation = 500 pg/ml)

If prostaglandin E₂ secretion stimulation is induced by the non-lymphokine parathyroid hormone (10⁻⁸ molar) then a small inhibition of prostaglandin synthesis is still observed.

Concentration of Interleukin-6 (Units/ml)	Prostaglandin E ₂ pg/ml
0	3700
10	3450
100	3300

Interleukin-6 has no effect on prostaglandin E₂ synthesis of unstimulated bone cultures in the above test.

Test 3

IL-6 molecules increase bone formation as indicated by e.g. an increased incorporation of [3H]proline into collagen and non-collagen proteins in fetal rat calvaria.

Organ cultures of rat calvaria are prepared by dissecting frontal and parietal bones from 21-day old fetal rats, splitting along the sagittal suture and culturing according to the method of Kream et al. (Endocrinology (1985) 116, 296). The IL-6 molecules are added in doses of from 1 to 100 units to 2 ml of cultures. Culturing is effected for 24 to 48 hours.

To quantitate the incorporation of [3H]proline into collagenase-digestible protein and non-collagen protein, bone homogenates are digested with bacterial collagenase according to the method of Diegelmann R. and Peterkofsky (Dev.Biol. (1972) 28:443) and modified by Kream et al. (Endocrinology (1985) 116:296).

Test 4

11-6 molecules decrease bone resorption as indicated by a decrease in release of [45]Ca from bone.

The test is effected according to the principles of Raisz (J.Clin.Invest. (1965) 44, 103).

- 5 Pregnant rats are injected s.c. with [45]Ca on the eighteenth day of gestation. Placebo or IL-6 molecules (1 to 100 Units per animal) are injected s.c. On day nineteen, the animals are sacrificed, the fetuses removed. The mineralized shafts of the radii and ulnae are dissected and placed in culture. Resorption is quantitated on the basis of release of [45]Ca from the bone explants.

Test 5

- 10 IL-6 molecules decrease bone resorption as indicated by a decreased number of osteoclasts and an increased amount of bone as indicated in the following test:-

- 15 Growing mice (6 to 8 weeks old; weight ca. 20 grams) are used. IL-6 molecules in doses of from about 0.1 to about 10 micrograms, e.g. 1 microgram, are injected subcutaneously over the calvarial bones, e.g. over the frontal and temporal bones of the skull. Injections are made daily for three to ten days. 3 days, 7 days or 2 weeks after the cessation of treatment the animals are sacrificed. The calvarial bones are examined by histology. For example the frontal and temporal bones are fixed, embedded and sectioned. The decrease in bone resorption relative to control animals is observed in a morphological basis, e.g. by more bone matrix and by a decreased number of osteoclasts as revealed by staining e.g. with tartrate resistant acid phosphatase.

- 20 Moreover, an increase in bone formation in the calvarial bones, e.g. frontal and temporal bones, relative to control animals is observed on a morphological basis, e.g. by an increased bone growth, and also observed by an increase in the number of osteoblasts.

Test 6

- 25 Release of Interleukin-6 like activity is stimulated in human osteoblast-like Saos-2 cells by phorbol 12-myristate 13-acetate (PMA) (which is a known stimulator of interleukin-6 from non-bone cell cultures) (at a concentration 2 to 200 of nanogram/ml), interleukin-1 (1 to 20 units/ml) and parathyroid hormone (PTH) (10^{-7} to 10^{-10} M).

- 30 In this test the osteoblastic human osteosarcoma cell line Saos-2 is cultured in McCoy's 5A medium supplemented 15% (v/v) fetal calf serum (FCS), penicillin, streptomycin and 2 mM L-glutamine in a humidified atmosphere of 5% CO₂ in air at 37°C. At confluency, cells are refed with medium containing 2 or 10% fetal calf serum and supplemented with the different agents to be tested. The medium of the Saos-2 cells is harvested after a 48 hour or 72 hour stimulation period and stored at 4°C.

- 35 Interleukin-6 like activity is detected using a conventional tritium -thymidine incorporation assay as follows:- Conditioned medium of Soas-2 cells are assayed for the presence of interleukin-6 activity using the above-mentioned B-cell (clone B13.29) proliferation assay. Significant levels of Interleukin-6 are detected within 48 hours.

Representative results are as follows:-

FCS = 2%

Dilution of conditional medium 1/10

Stimulator	³ H-thymidine incorporation (cpm)*
None	874
45 PMA (20 mg/ml)	61015
IL-1 (10 U/ml)	23189
PTH (10^{-8} M)	1287

- 50 * (background = 400 cpm; maximum incorporation 66000 cpm)

No detectable amounts of interleukin-6 like activity under control non-stimulated conditions are observed.

Test 7

- 55 Culturing of the mouse calvarial cells described above under test 2 gives significant interleukin-6 activity under control conditions (20000 cpm) after 48 hours culture. This is stimulated in the presence of phorbol 12-myristate 13-acetate (PMA) (1 to 100 ng /ml) and parathyroid hormone, hPTH₁₋₃₄, (PTH) (10^{-7} to 10^{-10} M).

Results obtained (³H thymidine incorporation in cpm) are:-

60

65

	Dilution	Control	PMA 100 ng/ml	PMA 10 ng/ml	PMA 1 ng/ml	
1	10	39919	47510	37679	48150	
2	100	9075	48239	36162	18145	
3	1000	680	27694	17687	1370	5
4	10000	346	5885	2545	580	
5	100000	305	595	572	411	
<hr/>						
	PTH 10-7 M	7 PTH 10-8 M	8 PTH 10-9 M	9 PTH 10-10 M		
1	12326	14638	16252	35897		10
2	23222	26885	30172	13073		
3	22816	18503	9059	827		
4	1935	1279	1095	531		
5	336	452	682	678		15

The present invention provides accordingly

- i) a method of inhibiting bone resorption or increasing bone formation or treating osteoporosis or osteoarthritis or any other indication mentioned above in a subject which comprises administering a therapeutically effective amount of an IL-6 molecule to a subject in need of such treatment, 20
- ii) the use of an IL-6 molecule in the manufacture of a medicament suitable for inhibiting bone resorption or increasing bone formation or treating osteoporosis or osteoarthritis or any other indication mentioned above,
- iii) the use of an IL-6 molecule for inhibiting bone resorption or increasing bone formation or treating osteoporosis or osteoarthritis or any other indication mentioned above, or 25
- iv) a medicament for inhibiting bone resorption or increasing bone formation or treating osteoporosis or osteoarthritis or any other indication mentioned above containing an IL-6 molecule. By IL-6 molecule is meant any compound corresponding to, having essentially the same structure as, or the same spectrum of activity as the known interleukin-6 (see the above mentioned PCT application or in the specific proliferation assay referred to above with reference to the Landsdorp and Aarden publications) or factors mentioned above, e.g. interferon beta-2 as described in e.g. EP 220574A (Yeda), PCT application WO 88/00206 (Genetics Institute), and EP 257406 (Ajinomoto). The contents of all these publications (hereinafter interleukin-6 publications) are incorporated herein by reference. 30

The IL-6 molecule may be glycosylated, e.g. on the arginine sites, and may be produced, e.g. from E.coli or CHO cells. 35

The IL-6 molecule as described is a 212 amino acid polypeptide. If desired the first or last methionine amino acid or even the whole non-coding region may be omitted, i.e. the first 28 amino acids, from Met.Asn up to and including Phe Pro Ala.

The interleukin-6 may have one, two or three or more amino acids absent, added or replaced by others and still retain interleukin-6 like activity, and be active in the method of the invention. 40

The expression IL-6 molecule covers such analogues as well as any compound which selectively stimulates IL-6 activity in bone cultures, especially if the compound is selective in that it has an immaterial action on other factors or bone resorption by other routes.

The term thus includes a compound of formula I:- 45

50

55

60

65

PRO VAL PRO PRO GLY GLU ASP SER LYS ASP VAL
 ALA ALA PRO HIS ARG GLN PRO LEU THR SER SER
 5 GLU ARG ILE ASP LYS GLN ILE ARG TYR ILE LEU
 ASP GLY ILE SER ALA LEU ARG LYS GLU THR CYS
 ASN LYS SER ASN MET CYS GLU SER SER LYS GLU
 10 ALA LEU ALA GLU ASN ASN LEU ASN LEU PRO LYS
 MET ALA GLU LYS ASP GLY CYS PHE GLN SER GLY
 PHE ASN GLU GLU THR CYS LEU VAL LYS ILE ILE
 15 THR GLY LEU LEU GLU PHE GLU VAL TYR LEU GLU
 TYR LEU GLN ASN ARG PHE GLU SER SER GLU GLU
 GLN ALA ARG ALA VAL GLN MET SER THR LYS VAL
 20 LEU ILE GLN PHE LEU GLN LYS LYS ALA LYS ASN
 LEU ASP ALA ILE THR THR PRO ASP PRO THR THR
 ASN ALA SER LEU LEU THR LYS LEU GLN ALA GLN
 25 ASN GLN TRP LEU GLN ASP MET THR THR HIS LEU
 ILE LEU ARG SER PHE LYS GLU PHE LEU GLN SER
 SER LEU ARG ALA LEU ARG GLN MET.

30

It also includes a compound of formula I having additional amino acids, e.g. attached to the N-terminal group, e.g. Ala or Met Ala or of formula II.

35

ALA PRO THR SER SER SER THR LYS LYS THR GLN
 LEU GLN LEU GLU HIS LEU LEU LEU ASP LEU PHE (II)
 40 ARG ALA - X

wherein X is formula I.

45 The term also includes molecules wherein one or more amino acids are replaced or omitted, e.g. incorporating the sequence.

50 PRO VAL PRO PRO GLY GLU ASP SER LYS
 ASP VAL ALA ALA

e.g. as the N-terminal sequence, and/or retaining essentially the same sequence of the remainder.

55 For these indications, the appropriate dosage will, of course, vary depending upon, for example, exact IL-6 molecule employed, the host, the mode of administration and the nature and severity of the condition being treated. However, in general, satisfactory results in animals are indicated to be obtained at daily dosages from about 0.5 micrograms/kg to about 20 micrograms/kg animal body weight. In larger mammals, for example humans, an indicated daily dosage is in the range from about 25 micrograms to about 1000 micrograms of an
 60 IL-6 molecule conveniently administered, for example, in divided doses up to four times a day.

The IL-6 molecules may be administered by any conventional route, in particular enterally e.g. in the form of tablets or capsules, or preferably parenterally, e.g. in the form of injectable solutions or suspensions.

For example methods of administration and galenical formulations for use interleukin-6 are known, e.g. from the above-mentioned interleukin-6 publications.

65 Interleukin-6 is the preferred compound. Test results are given above. It is indicated that the compound may

be administered at daily dosages of from 25 micrograms to 125 micrograms by subcutaneous injection to larger mammals, for example humans.

Pharmaceutical compositions may comprise an IL-6 molecule in association with at least one pharmaceutical carrier or diluent. Such compositions may be manufactured in conventional manner. Unit dosage forms contain, for example, from about 4 micrograms to about 500 micrograms of the IL-6 molecule.

The present composition also provides a pack containing a pharmaceutical composition comprising an IL-6 molecule together with instructions for use in a method as defined above.

Claims

1. The use of an IL-6 molecule in the manufacture of a medicament suitable for inhibiting bone resorption.
2. The use of an IL-6 molecule in the manufacture of a medicament suitable for increasing bone formation.
3. The use of an IL-6 molecule in the manufacture of a medicament suitable for treating osteoporosis.
4. The use of an IL-6 molecule in the manufacture of a medicament suitable for treating osteoarthritis.
5. The use of an IL-6 molecule in the manufacture of a medicament suitable for treating at least one of the following diseases; Paget's disease of bone, osteolysis due to bone malignancies, osteomalacia, osteogenesis imperfecta and osteomyelitis.
6. The use of an IL-6 molecule in the manufacture of a medicament suitable for treating multiple myeloma of the bone.
7. The use of an IL-6 molecule in the manufacture of a medicament suitable for treating hypercalcemia.
8. A use according to any preceeding claim wherein the IL-6 molecule is interleukin-6.
9. A use according to any preceeding claim wherein the IL-6 molecule is of formula I:-

PRO VAL PRO PRO GLY GLU ASP SER LYS ASP VAL
 ALA ALA PRO HIS ARG GLN PRO LEU THR SER SER
 GLU ARG ILE ASP LYS GLN ILE ARG TYR ILE LEU
 ASP GLY ILE SER ALA LEU ARG LYS GLU THR CYS
 ASN LYS SER ASN MET CYS GLU SER SER LYS GLU
 ALA LEU ALA GLU ASN ASN LEU ASN LEU PRO LYS
 MET ALA GLU LYS ASP GLY CYS PHE GLN SER GLY
 PHE ASN GLU GLU THR CYS LEU VAL LYS ILE ILE
 THR GLY LEU LEU GLU PHE GLU VAL TYR LEU GLU
 TYR LEU GLN ASN ARG PHE GLU SER SER GLU GLU
 GLN ALA ARG ALA VAL GLN MET SER THR LYS VAL
 LEU ILE GLN PHE LEU GLN LYS LYS ALA LYS ASN
 LEU ASP ALA ILE THR THR PRO ASP PRO THR THR
 ASN ALA SER LEU LEU THR LYS LEU GLN ALA GLN
 ASN GLN TRP LEU GLN ASP MET THR THR HIS LEU
 ILE LEU ARG SER PHE LYS GLU PHE LEU GLN SER
 SER LEU ARG ALA LEU ARG GLN MET.

I

10. A use according to claim 8 wherein the compound is Ala-X wherein X is the sequence of formula I.

11. A use according to claim 8 wherein the compound is Met-Ala-X wherein X is the sequence of formula I.
12. A use according to claim 9 wherein the IL-6 molecule incorporates the sequence

5

PRO VAL PRO PRO GLY GLU ASP SER LYS
ASP VAL ALA ALA.

10

13. A use according to claim 1 substantially as hereinbefore described with reference to any one of the examples.

14. A method of treating one of the diseases mentioned in any preceeding claim which comprises administering a compound as defined in any preceeding claim to a subject in need of such treatment.

15

15. A pack containing a pharmaceutical composition comprising an IL-6 molecule together with instructions for use in a method as defined in claim 14.

20

25

30

35

40

45

50

55

60

65

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 331 640 A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **89810155.5**

(51) Int. Cl.⁵: **A61K 37/02**

(22) Date of filing: **28.02.89**

(30) Priority: **04.03.88 GB 8805231**
09.05.88 GB 8810899

(43) Date of publication of application:
06.09.89 Bulletin 89/36

(64) Designated Contracting States:
ES GR LU

(68) Date of deferred publication of the search report:
13.03.91 Bulletin 91/11

(71) Applicant: **SANDOZ AG**
Lichtstrasse 35
CH-4002 Basel(CH)

(72) Inventor: **Feyen, Jean Honoré M.**
Mühlegasse 2
CH-4104 Oberwil(CH)
Inventor: **Trechsel, Ulrich**
Bollwerkstrasse 60
CH-4102 Binningen(CH)

(54) **Use of an interleukin-6 molecule in inhibiting bone resorption.**

(57) We have found that IL-6 molecules are useful in inhibiting bone resorption and increasing bone formation. They are thus useful in treating osteoporosis, e.g. acute and/or chronic stages of postmenopausal osteoporosis, including primary and/or secondary osteoporosis and localized osteoporosis, post-traumatic osteoporosis, Paget's disease of bone (osteitis deformans), osteolysis due to bone malignancies, e.g. osteogenic sarcoma, osteodystrophy, e.g. renal osteodystrophy, osteomalacia, multiple myeloma of the bone, osteogenesis imperfecta, hypercalcemia, osteomyelitis, e.g. in conjunction with antibiotics and ectopic calcification.

EP 0 331 640 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

Application number

EP 89 81 0155

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A,D	EP-A-0 257 406 (AJINMOTO) * Claims 1-15,46-47; page 8, line 55 - page 13, line 8 * --	1-12, 15	A 61 J 37/02
A	J. EXP. MED., vol. 165, March 1987, pages 914-919, The Rockefeller University Press; J. VAN DAMME et al.: "Identification of the human 26kD protein, interferon beta 2 (IFN-beta 2), as A B cell hybridoma/plasmacytoma growth factor induced by interleukin I and tumor necrosis factor" * Page 918, Summary; page 914, Introduction * -- ./.	1-12, 15	TECHNICAL FIELDS SEARCHED (Int. Cl.4) A 61 K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-7 Claims searched incompletely: 13-14 Claims not searched: 13-14 Reason for the limitation of the search:</p> <p>Claims 13-14: EPC Rule 29(6) and Method for treatment of the human or animal body by surgery or therapy (See art. 52(4) of the European Patent Convention) Claims 1-7: The term "a IL-6 molecule" (claims 1-7) doesn't make completely clear, which compound is meant. EPC Art. 84.</p>			
Place of search THE HAGUE		Date of completion of the search 17-11-1990	Examiner ISERT
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application number

EP 89 81 0155

-2-

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	INT. J. CANCER, suppl. 1, 1987, pages 14-20, Alan R. Liss, Inc.; R. OHNO: "Interferons in the treatment of multiple myeloma" * Summary, page 17, table IV and lowest paragraph * -----	6	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)